Zenith[®] Fenestrated

AAA ENDOVASCULAR GRAFT

PHYSICIAN DICTATION NOTES

Date:	Hospital:				
Physician:			Attending physician:		
Patient ID:	Tin	ne of procedure:	Fluoroscopy time:	Total contrast:	Ipsilateral: 🗆 RT 🗅 LT
Devices used					
Proximal body graft:			Additional devices:		
Distal body graft:					
Controlatoral iliac logi					

Coding information*


34831

+37237

endovascular repair; aorto-bi-iliac prosthesis

each additional artery

(superior mesenteric, celiac and/or renal artery[s])



Comments:



penetrating ulcer, intramural hematoma, or traumatic disruption) with a fenestrated visceral aortic endograft and concomitant unibody or modular infrarenal aortic endograft and all associated radiological supervision and interpretation, including target zone angioplasty, when performed; including three visceral artery endoprostheses (superior mesenteric, celiac and/or renal artery[s]) Endovascular repair of visceral aorta and infrarenal abdominal aorta (eg, aneurysm, pseudoaneurysm, dissection, penetrating ulcer, intramural hematoma, or traumatic disruption) with a fenestrated visceral aortic endograft and concomitant unibody or modular infrarenal aortic endograft and all associated radiological supervision and interpretation, including target zone angioplasty, when performed; including four or more visceral artery endoprostheses (superior mesenteric, celiac and/or renal artery[s]) Placement of extension prosthesis(es) distal to the common iliac artery(ies) or proximal to the renal artery(ies) for endovascular repair of infrarenal abdominal aortic or iliac aneurysm, false aneurysm, dissection, penetrating ulcer, including pre-procedure sizing and device selection, all nonselective catheterization(s), all associated radiological supervision and interpretation, and treatment zone angiopasty/stenting, when performed, per vessel treated (34709 may only be reported once per vessel treated [ie, multiple endograft extensions placed in a single vessel may only Percutaneous access and closure of femoral artery for delivery of endograft through a large sheath (12French or larger), including ultrasound guidance, when performed, unilateral (34713 may only be reported once per side. For bilateral procedure, report 34713 twice) (Do not report ultrasound guidance [ie, 76937] for percutaneous vascular access in conjunction with 34713 for the same access +34812 Open femoral artery exposure for delivery of endovascular prosthesis, by groin incision, unilateral (34812 may only be reported once per side. For bilateral procedure, report 34812 twice) Open femoral artery exposure with creation of conduit for delivery of endovascular prosthesis or for establishment +34714 of cardiopulmonary bypass, by groin incision, unilateral (34714 may only be reported once per side. For bilateral procedure, report 34714 twice) Open iliac artery exposure for delivery of endovascular prosthesis or iliac occlusion during endovascular therapy, by +34820 abdominal or retroperitoneal incision, unilateral (34820 may only be reported once per side. For bilateral procedure, report 34820 twice) Open iliac artery exposure with creation of conduit for delivery of endovascular prosthesis or for establishment of cardiopulmonary bypass, by abdominal or retroperitoneal incision, unilateral (34833 may only be reported once per side. For bilateral procedure, report 34833 twice) +34834 Open brachial artery exposure for delivery of endovascular prosthesis, unilateral (34834 may only be reported once per side. For bilateral procedure, report 34834 twice) +34715 Open axillary/subclavian artery exposure for delivery of endovascular prosthesis by infraclavicular or supraclavicular inicision, unilateral (34715 may only be reported once per side. For bilateral procedure, report 34715 twice) Open axillary/subclavian artery exposure with creation of conduit for delivery of endovascular pros +34716 establishment of cardiopulmonary bypass, by infraclavicular or supraclavicular incision, unilateral (34716 may only be reported once per side. For bilateral procedure, report 34716 twice) +34813 Placement of femoral-femoral prosthetic graft during endovascular aortic aneurysm repair (Use 34813 in conjunction with 34812) 34830 Open repair of infrarenal aortic aneurysm or dissection, plus repair of associated arterial trauma, following unsuccessful endovascular repair; tube prosthesis Open repair of infrarenal aortic aneurysm or dissection, plus repair of associated arterial trauma, following unsuccessful

34839 Physician planning of a patient-specific fenestrated visceral aortic endograft requiring a minimum of 90 minutes of physician time Endovascular repair of visceral aorta and infrarenal abdominal aorta (eg. aneurysm, pseudoaneurysm, dissection, penetrating ulcer, intramural hematoma, or traumatic disruption) with a fenestrated visceral aortic endograft and 34845 concomitant unibody or modular infrarenal aortic endograft and all associated radiological supervision and $interpretation, including \ target \ zone \ angioplasty, when \ performed; including \ one \ visceral \ artery \ endoprost hesis \ (superior \ mesenteric, \ celiac \ or \ renal \ artery)$

Endovascular repair of visceral aorta and infrarenal abdominal aorta (eg. aneurysm, pseudoaneurysm, dissection, penetrating ulcer, intramural hematoma, or traumatic disruption) with a fenestrated visceral aortic endograft and concomitant unibody or modular infrarenal aortic endograft and all associated radiological supervision and interpretation, including target zone angioplasty, when performed; including two visceral artery endoprostheses

Endovascular repair of visceral aorta and infrarenal abdominal aorta (eg, aneurysm, pseudoaneurysm, dissection,

*This information should not be construed as authoritative. The entity billing Medicare and/or third party payers is solely responsible for codes assigned to the services and items in the medical record. is solely responsible for codes assigned to the services and items in the mo

34832 Open repair of infrarenal aortic aneurysm or dissection, plus repair of associated arterial trauma, following unsuccessful endovascular repair; aorto-bifemoral prosthesis Transcatheter placement of an intravascular stent(s) (except lower extremity artey(s) for occlusive disease, cervical carotid,

37236 extracranial vertebral or intrathoracic carotid, intracranial, or coronary), open or percutaneous, including radiological supervision and interpretation and including all angioplasty within the same vessel, when performed; initial artery Transcatheter placement of an intravascular stent(s) (except lower extremity artev(s) for occlusive disease, cervical carotid, extracranial vertebral or intrathoracic carotid, intracranial, or coronary), open or percutaneous, including radiological supervision and interpretation and including all angioplasty within the same vessel, when performed;

CPT© 2017 American Medical Association. All rights reserved. CPT is a registered trademark of the American Medical Association